

Returns to the Origin for Random Walks on \mathbb{Z} Revisited

Helmut Prodinger

The John Knopfmacher Centre for Applicable Analysis and Number Theory

School of Mathematics, University of the Witwatersrand, P. O. Wits

2050 Johannesburg, South Africa

helmut@maths.wits.ac.za

Chrysafi and Bradley [1] consider symmetric random walks, defined as follows: Let X_k , $k = 1, 2, \dots$ be independent and identically distributed random variables with $\mathbb{P}\{X_k = 1\} = \mathbb{P}\{X_k = -1\} = \frac{1}{2}$. Then

$$S_m = \sum_{k=1}^m X_k \quad \text{with} \quad S_0 = 0$$

is a simple random walk starting at 0. The authors considered only walks of even length $m = 2n$ and were interested in the random variable $R = R_n$, defined to be the NUMBER OF RETURNS TO THE ORIGIN in a walk of length $2n$, i.e., the number of times $S_i = 0$ happens, for $i = 1, \dots, 2n$. They computed moments up to $\mathbb{E}[R^6]$ and ask for a closed formula for $\mathbb{E}[R^k]$ and also whether $\mathbb{E}[R^k] \sim c_k n^{k/2}$ holds.

The answers to these questions can be found in [4], compare also [5]. There, the *factorial moments* $\mathbb{E}[R^{\underline{k}}]$ were computed. We state the formula only for even n :

$$\mathbb{E}[R_n^{\underline{k}}] = k! \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} \binom{\frac{i}{2} + n}{n}.$$

Ordinary moments can be recovered from these formulæ as linear combinations with *Stirling numbers of the second kind* (*Stirling subset numbers*), see [3]:

$$\mathbb{E}[R_n^k] = \sum_{i=0}^k \left\{ \begin{matrix} k \\ i \end{matrix} \right\} \mathbb{E}[R_n^{\underline{i}}].$$

To answer the asymptotics question, we use generating functions:

$$\mathbb{E}[R_n^k] = 4^{-n} k! [z^n] \frac{(1 - \sqrt{1 - 4z})^k}{(1 - 4z)^{1 + \frac{k}{2}}},$$

and *singularity analysis of generating functions*, as nicely described in [2]. One must expand around the dominant singularity (here $z = \frac{1}{4}$):

$$\frac{(1 - \sqrt{1 - 4z})^k}{(1 - 4z)^{1 + \frac{k}{2}}} \sim (1 - 4z)^{-1 - \frac{k}{2}}$$

and use a transfer theorem:

$$\mathbb{E}[R_n^k] \sim 4^{-n} k! [z^n] (1 - 4z)^{-1 - \frac{k}{2}} = k! \binom{\frac{k}{2} + n}{n} \sim \frac{k!}{(k/2)!} n^{k/2}.$$

For odd $k = 2j + 1$, the factor may be rewritten as follows:

$$\frac{k!}{(k/2)!} = \frac{(2j + 1)!}{(j + \frac{1}{2})!} = \frac{2^{2j+1} j!}{\sqrt{\pi}}.$$

For ordinary moments, the leading terms in the asymptotic expansion are the same:

$$\mathbb{E}[R_n^k] \sim \frac{k!}{(k/2)!} n^{k/2}.$$

REFERENCES

- [1] L. A. Chrysafti and R. E. Bradley. Returns to the origin for random walks on \mathbb{Z} . *Missouri J. Math. Sci.*, 14(2):96–106, 2002.
- [2] P. Flajolet and A. Odlyzko. Singularity analysis of generating functions. *SIAM Journal on Discrete Mathematics*, 3:216–240, 1990.
- [3] R. L. Graham, D. E. Knuth, and O. Patashnik. *Concrete Mathematics: a foundation for computer science*. Addison-Wesley Publishing Company, Amsterdam, 2nd edition, 1994.
- [4] P. Kirschenhofer and H. Prodinger. The higher moments of the number of returns of a simple random walk. *Adv. in Appl. Probab.*, 26(2):561–563, 1994.
- [5] P. Kirschenhofer and H. Prodinger. Return statistics of simple random walks. *J. Statist. Plann. Inference*, 54(1):67–74, 1996.