

# COMPUTER-FREE EVALUATION OF A DOUBLE INFINITE SUM VIA EULER SUMS

ALOIS PANHOLZER AND HELMUT PRODINGER

ABSTRACT. For a double sum that was recently computed by Pemantle and Schneider using the computer software *Sigma*, a short and computer-free proof is provided.

## 1. INTRODUCTION

The evaluation of the double sum

$$S := \sum_{j,k=1}^{\infty} \frac{H_j(H_{k+1} - 1)}{jk(k+1)(j+k)} = -\zeta(2) - 2\zeta(3) + 4\zeta(2)\zeta(3) + 2\zeta(5), \quad (1)$$

(where  $H_n := \sum_{k=1}^n \frac{1}{k}$  denote harmonic numbers) appears in [4] and was obtained using Carsten Schneider's software *Sigma*. Here, we will give a short proof that is completely computer-free.

The second order harmonic numbers which appear in the sequel are denoted by  $H_n^{(2)} := \sum_{k=1}^n \frac{1}{k^2}$ .

We split the sum  $S$  and apply partial fraction decomposition:

$$S = \sum_{k \geq 1} \frac{H_{k+1} - 1}{k(k+1)} \sum_{j \geq 1} \frac{H_j}{j(j+k)} = \sum_{k \geq 1} \frac{H_{k+1} - 1}{k^2(k+1)} \sum_{j \geq 1} H_j \left( \frac{1}{j} - \frac{1}{j+k} \right).$$

The inner sum is simplified as follows:

$$\sum_{j \geq 1} H_j \left( \frac{1}{j} - \frac{1}{j+k} \right) = \sum_{j \geq 1} \left( \frac{1}{j} - \frac{1}{j+k} \right) \sum_{l=1}^j \frac{1}{l} = \sum_{l \geq 1} \frac{1}{l} \sum_{j \geq l} \left( \frac{1}{j} - \frac{1}{j+k} \right).$$

The second sum here telescopes, and only  $k$  summands remain:

$$\begin{aligned} \sum_{l \geq 1} \frac{1}{l} \sum_{j \geq l} \left( \frac{1}{j} - \frac{1}{j+k} \right) &= \sum_{l \geq 1} \frac{1}{l} \sum_{j=0}^{k-1} \frac{1}{l+j} = \sum_{l \geq 1} \frac{1}{l^2} + \sum_{j=1}^{k-1} \sum_{l \geq 1} \frac{1}{l(l+j)} \\ &= \zeta(2) + \sum_{j=1}^{k-1} \frac{1}{j} \sum_{l \geq 1} \left( \frac{1}{l} - \frac{1}{l+j} \right) \quad [\text{partial fraction decomposition}] \\ &= \zeta(2) + \sum_{j=1}^{k-1} \frac{H_j}{j} \quad [\text{again by telescoping}] \\ &= \zeta(2) + \frac{H_k^2 + H_k^{(2)}}{2} - \frac{H_k}{k}. \end{aligned}$$

---

*Date:* January 20, 2005.

Thus the task reduces to evaluate the following single sum:

$$S = \sum_{k=1}^{\infty} \frac{H_{k+1} - 1}{k^2(k+1)} \left( \zeta(2) + \frac{H_k^2 + H_k^{(2)}}{2} - \frac{H_k}{k} \right). \quad (2)$$

After partial fraction expansion (and shifting the index if necessary), sum  $S$  can be written as follows:

$$S = -2\zeta(2) + \frac{1}{2} \sum_{k \geq 1} \frac{H_k H_k^{(2)}}{k^2} + \frac{1}{2} \sum_{k \geq 1} \frac{H_k^3}{k^2} - \sum_{k \geq 1} \frac{H_k^2}{k^3} + (\zeta(2) - 1) \sum_{k \geq 1} \frac{H_k}{k^2} - 2 \sum_{k \geq 1} \frac{1}{k^2}.$$

For the final computation of  $S$  we require the following evaluations of Euler sums via values of the Riemann zeta function:

$$\sum_{k \geq 1} \frac{H_k H_k^{(2)}}{k^2} = \zeta(5) + \zeta(2)\zeta(3), \quad (3a)$$

$$\sum_{k \geq 1} \frac{H_k^3}{k^2} = 10\zeta(5) + \zeta(2)\zeta(3), \quad (3b)$$

$$\sum_{k \geq 1} \frac{H_k^2}{k^3} = \frac{7}{2}\zeta(5) - \zeta(2)\zeta(3), \quad (3c)$$

$$\sum_{k \geq 1} \frac{H_k}{k^2} = 2\zeta(3), \quad (3d)$$

from which equation (1) then follows.

Equations (3c) and (3d) can be found explicitly in [3]. In [3] one also finds the identities

$$\sum_{k \geq 1} \frac{H_k^3}{(k+1)^2} = \frac{15}{2}\zeta(5) + \zeta(2)\zeta(3), \quad (4a)$$

$$\sum_{k \geq 1} \frac{H_k}{k^4} = 3\zeta(5) - \zeta(2)\zeta(3), \quad (4b)$$

and due to

$$\sum_{k \geq 1} \frac{H_k^3}{k^2} = \sum_{k \geq 1} \frac{H_k^3}{(k+1)^2} + 3 \sum_{k \geq 1} \frac{H_k^2}{k^3} - 3 \sum_{k \geq 1} \frac{H_k}{k^4} + \sum_{k \geq 1} \frac{1}{k^5},$$

we obtain equation (3b) as well, using (4a), (4b) and (3c).

To show (3a) we will apply Theorem 2 of [2], which gives

$$\sum_{k \geq 1} \frac{H_{k-1} H_{k-1}^{(2)}}{k^2} = \zeta(2, 1, 2) + \zeta(2, 2, 1) + \zeta(2, 3),$$

where the *multiple zeta functions* are defined by

$$\zeta(a_1, a_2, \dots, a_m) = \sum_{k_1 > k_2 > \dots > k_m \geq 1} \frac{1}{k_1^{a_1} k_2^{a_2} \dots k_m^{a_m}}.$$

Since

$$\begin{aligned} \sum_{k \geq 1} \frac{H_k H_k^{(2)}}{k^2} &= \sum_{k \geq 1} \frac{H_{k-1} H_{k-1}^{(2)}}{k^2} + \sum_{k \geq 1} \frac{H_{k-1}^{(2)}}{k^3} + \sum_{k \geq 1} \frac{H_{k-1}}{k^4} + \sum_{k \geq 1} \frac{1}{k^5} \\ &= \sum_{k \geq 1} \frac{H_{k-1} H_{k-1}^{(2)}}{k^2} + \sum_{k > l \geq 1} \frac{1}{k^3 l^2} + \sum_{k > l \geq 1} \frac{1}{k^4 l} + \sum_{k \geq 1} \frac{1}{k^5}, \end{aligned}$$

we obtain

$$\sum_{k \geq 1} \frac{H_k H_k^{(2)}}{k^2} = \zeta(2, 1, 2) + \zeta(2, 2, 1) + \zeta(2, 3) + \zeta(3, 2) + \zeta(4, 1) + \zeta(5).$$

Using the following evaluations of the multiple zeta function given in [1] resp. [2]:

$$\begin{aligned} \zeta(2, 1, 2) &= \zeta(2, 3) = \frac{9}{2} \zeta(5) - 2\zeta(2)\zeta(3), \\ \zeta(2, 2, 1) &= \zeta(3, 2) = -\frac{11}{2} + 3\zeta(2)\zeta(3), \\ \zeta(4, 1) &= 2\zeta(5) - \zeta(2)\zeta(3), \end{aligned}$$

equation (3a) follows.

#### REFERENCES

- [1] D. Borwein, J. Borwein and R. Girgensohn, Explicit evaluation of Euler sums, *Proceedings of the Edinburgh Mathematical Society* 38, 277–294, 1995.
- [2] J. Borwein and R. Girgensohn, Evaluation of triple Euler sums, *Electronic Journal of Combinatorics* 3, research paper #23, 1996.
- [3] P. Flajolet and B. Salvy, Euler sums and contour integral representations, *Experimental Mathematics* 7, 15–35, 1998.
- [4] R. Pemantle and C. Schneider, When is 0.999... equal to 1?, preprint, 2004.

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNIVERSITÄT WIEN, WIEDNER HAUPTSTRASSE 8–10/104, A-1040 WIEN, AUSTRIA.

*E-mail address:* Alois.Panholzer@tuwien.ac.at

HELMUT PRODINGER, MATHEMATICS DEPARTMENT, STELLENBOSCH UNIVERSITY, 7602 STELLENBOSCH, SOUTH AFRICA.

*E-mail address:* hproding@sun.ac.za